

NAVAL POSTGRADUATE SCHOOL
Monterey, California

EC 3210

FINAL EXAM

12/89Po

- This exam is open book and notes.
- There are five problems; each is equally weighted.
- Partial credit will be given; be sure to do some work on each problem.
- Be sure to include units in your answers.
- Please circle or underline your answers.
- Show *ALL* work.
- Write only your name on this sheet.
- Exams and course grades will be available in the ECE Office in Sp 437 on Tuesday, 26 December.

Course grade: _____

1		4	
2		5	
3			
Total			

Name: _____

1. The carbon-monoxide (CO) laser is a four-level laser whose energy levels have the properties listed below. The laser operates at a cryogenic temperature of 77K at a pressure of 10 Torr. Write an expression for the linewidth function $g(\nu)$ if the laser is lifetime and collision broadened. (The collision cross-section for a CO-CO collision is $7.03 \times 10^{-21} \text{ m}^{-2}$.)

Energy level	Energy (eV)	Lifetime (μs)
3	3.10	1.0×10^{-3}
2	1.91	2.0
1	1.5	0.10
Ground	0	∞

(Note: This data is fictitious and does not represent the properties of an actual CO laser.)

2. A Gaussian laser beam ($\lambda = 800 \text{ nm}$) propagates from left to right. The beam has a spot size of 8 mm and a radius of curvature of phase of +15 m at a location that is 25 m from the origin (of the $x - y - z$ coordinate system). Find the beam spot size *and* radius of curvature of the phase at a location that is 9.5 m from the origin.
3. A greybody emitter has an emissivity of 0.8 and a temperature of 500K.
 - (a) Calculate the wavelength of peak spectral photon emittance. (5 points)
 - (b) Suppose that the emissivity is reduced to a value that is 50% of its original value and that the temperature is raised to a value that is 110% of its initial value. Calculate the factor by which the spectral photon emittance will change at the wavelength calculated in Part a. (15 points)
4. A mode-locked laser produces a train of pulses having a peak power of 10 watts, an average power of 50 mW, and a period of 60 ps. Calculate the longitudinal coherence length of this laser.
5. Calculate the required unsaturated population inversion *density* (in units of $\text{atoms} \cdot \text{m}^{-3}$) to achieve an unsaturated gain coefficient of 0.5 m^{-1} in CO_2 . The transition wavelength is $10.6 \text{ } \mu\text{m}$ and the Einstein absorption coefficient is $A_{21} = 200 \text{ s}^{-1}$. The transition is Doppler-broadened with a temperature of 400K. You may assume that $n = 1$.